



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

SOLUTION BY A. H. WILSON, Haverford College.

The discussion is for the center of the hexagon, an obvious position of equilibrium. Let this point be taken as origin, the hexagon placed with two vertices on the  $x$ -axis. Let  $r_0$  be the side of the hexagon.

Form the potential function  $V$ . For one attracting center this is of the form  $\int_{r_0}^r r^{-h} dV = r^{1-h}/(1-h) - r_0^{1-h}/(1-h)$ . (The value  $h = 1$  must be excepted.) For simplicity the origin is taken as a standard position.

$$V = \Sigma r^{1-h}/(1-h) - V_0.$$

Each  $r$  is of the form  $[(x-s)^2 + (y-t)^2]^{1/2}$ . Expand  $V$  in the neighborhood of the origin in powers of  $x$  and  $y$ . For  $r^{1-h}$  this expansion is

$$\begin{aligned} r^{1-h} = & r_0^{1-h} + s(1-h)r_0^{-h-1}x + t(1-h)r_0^{-h-1}y + \frac{1}{2}[(1-h)r_0^{-h-1} - (1-h^2)s^2r_0^{-h-3}]x^2 \\ & - (1-h^2)r_0^{-h-3}st \cdot xy + \frac{1}{2}[(1-h)r_0^{-h-1} - (1-h^2)r_0^{-h-3}t^2]y^2 + \dots \end{aligned}$$

For  $(s, t)$  substitute successively the coördinates of the hexagon points and sum:

$$V = V_0 - \frac{1}{2}(2 + 3h)r_0^{-h-1}(x^2 + y^2) + \dots$$

In the neighborhood of the origin then the equations of motion are

$$\begin{aligned} m \frac{d^2x}{dt^2} &= - \frac{\partial V}{\partial x} = (2 + 3h)r_0^{-h-1}x, \\ m \frac{d^2y}{dt^2} &= - \frac{\partial V}{\partial y} = (2 + 3h)r_0^{-h-1}y. \end{aligned}$$

For stable equilibrium we must have here

$$2 + 3h < 0.$$

## MISCELLANEOUS QUESTIONS.

EDITED BY R. D. CARMICHAEL.

### NEW QUESTIONS.

19. How many known proofs are there of the proposition that the square of the hypotenuse of a right-angled triangle is equal to the sum of the squares of the other two sides? Where are the proofs to be found?

20. Some of our readers would like to have a simple account, without proofs, of just what has been accomplished toward the proof of the theorem that the equation  $x^n + y^n = z^n$  is impossible in integers when  $n > 2$ .

21. For the diophantine equation

$$x^2 - y^3 = 17$$

there are known the following solutions:

$$\begin{aligned}x &= 3, & 4, & 5, & 9, & 23, & 282, & 375, & 378661, \\y &= -2, & -1, & 2, & 4, & 5, & 43, & 52, & 5234.\end{aligned}$$

One of our readers, who supplied the foregoing facts, desires to know the answers to the following questions: Are there other solutions of the given diophantine equation? How may all the solutions of this equation be found by a systematic procedure?

### REPLY.

14. In the process of solving a certain physical problem Professor H. S. Uhler, of Yale University, was led to the definite integral

$$\int_0^a (a^2 - x^2) x dx \int_{a-x}^{a+x} \frac{e^{-cy}}{y} dy,$$

for which he found the value

$$\frac{1}{c^2} \left[ a^2 - \frac{3a}{c} + \frac{1}{c} \left( a + \frac{3}{2c} \right) (1 - e^{-2ac}) \right],$$

$a$  and  $c$  being positive constants. Professor Uhler would like to see how other persons attack the problem of evaluating this integral.

### I. REMARKS BY A. M. HARDING, University of Arkansas.

Let

$$u = \int_0^a (a^2 - x^2) x f(x) dx,$$

where

$$f(x) = \int_{a-x}^{a+x} \frac{e^{-cy}}{y} dy.$$

Integrating by parts we have

$$u = -\frac{(a^2 - x^2)^2}{4} f(x) \Big|_0^a + \frac{1}{4} \int_0^a (a^2 - x^2)^2 f'(x) dx;$$

or,

$$u = \frac{1}{4} \int_0^a (a^2 - x^2)^2 f'(x) dx.$$

Now,

$$f'(x) = \frac{e^{-c(a+x)}}{a+x} + \frac{e^{-c(a-x)}}{a-x} = \frac{e^{-ac}}{a^2 - x^2} [a(e^{cx} + e^{-cx}) + x(e^{cx} - e^{-cx})].$$

Therefore

$$\begin{aligned}u &= \frac{ae^{-ac}}{4} \int_0^a (a^2 - x^2)(e^{cx} + e^{-cx}) dx + \frac{e^{-ac}}{4} \int_0^a (a^2 x - x^3)(e^{cx} - e^{-cx}) dx \\&= \frac{ae^{-ac}}{4} u_1 + \frac{e^{-ac}}{4} u_2, \text{ say.}\end{aligned}$$

Repeated integration by parts gives

$$\begin{aligned}u_1 &= \frac{a^2 - x^2}{c} (e^{cx} - e^{-cx}) \Big|_0^a + \frac{2}{c} \int_0^a x(e^{cx} - e^{-cx}) dx \\&= \frac{2}{c^2} \left[ a(e^{ac} + e^{-ac}) - \frac{1}{c} (e^{ac} - e^{-ac}) \right].\end{aligned}$$

$$\begin{aligned}
u_2 &= \frac{a^2x - x^3}{c} (e^{cx} + e^{-cx}) \Big|_0^a - \frac{1}{c} \int_0^a (a^2 - 3x^2)(e^{cx} + e^{-cx}) dx \\
&= -\frac{1}{c^2} \left[ (a^2 - 3x^2)(e^{cx} - e^{-cx}) \Big|_0^a + 6 \int_0^a x(e^{cx} - e^{-cx}) dx \right] \\
&= -\frac{1}{c^2} \left[ -2a^2(e^{ac} - e^{-ac}) + \frac{6}{c} \left\{ a(e^{ac} + e^{-ac}) - \frac{1}{c}(e^{ac} - e^{-ac}) \right\} \right] \\
&= \frac{1}{c^2} \left[ 2a^2(e^{ac} - e^{-ac}) - \frac{6a}{c}(e^{ac} + e^{-ac}) + \frac{6}{c^2}(e^{ac} - e^{-ac}) \right].
\end{aligned}$$

Substituting the values of  $u_1$  and  $u_2$  in the expression above for  $u$  we obtain the desired result.

---

## NOTES AND NEWS.

EDITED BY W. DEW. CAIRNS.

"The propagation of electric waves in wireless telegraphy" is the title of a paper by Professor George R. Dean, of the Missouri School of Mines, in *The Electrician* for September 11, 1914.

Dr. L. L. DINES, formerly professor of mathematics at the University of Arizona, has been elected to an associate professorship at the University of Saskatchewan.

The Southwestern Section of the American Mathematical Society held its eighth regular meeting at the University of Nebraska on Saturday, November 28, 1914.

Mr. J. L. RILEY, formerly instructor in mathematics at the University of Oklahoma, has accepted a fellowship in Rice Institute, Houston, Texas, for the year 1914-15. Professor E. P. R. DUVAL has returned to his former position as associate professor of mathematics at the University of Oklahoma.

Dr. E. E. Whitford has been promoted to an assistant professorship of mathematics at the College of the City of New York.

Mr. Joel D. Eshleman, A.B., has been appointed instructor in mathematics in Adelbert College, Western Reserve University.

Dr. E. B. Stouffer, formerly instructor in mathematics in the University of Illinois, has accepted an assistant professorship in the University of Kansas.

Professor M. Frechet, University of Poitiers, France, who was to lecture at the University of Illinois during the present year, has joined the French army.